



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ARTICLE XVII.

Observations to determine the horizontal Magnetic Intensity and Dip at Louisville, Kentucky, and at Cincinnati, Ohio. By John Locke, M. D., Professor of Chemistry in the Medical College of Ohio. Read May 15, 1840.

THE following observations were made to determine the relative horizontal magnetic intensities at Louisville, Kentucky, and Cincinnati, Ohio. The apparatus used was that invented by Professor Bache, in which the needles are made to oscillate in a rarefied medium. Two of the needles used (Nos. 1 and 2) were of the model of those of Professor Hansteen; the third (No. 3) was flat, with lozenge points.

HORIZONTAL INTENSITY.

| <i>Cincinnati, March 7, 1840. Latitude 39° 06' N.; Longitude 84° 27' W.</i> | | | | | | |
|---|--|------------|-----------------------------|-----------------------------|---------------------|----------------------------|
| No. of Needle. | Time of Commencement. | Mean Temp. | Duration of 500 Vibrations. | Calculated duration at 60°. | Square of the Time. | Observations. |
| No. 1 | 11 ^h 10 ^m 05 ^s .6 | 56° 64' F. | 25.6666 Min. | 25.6777 Min. | 659.34428 | Cloudy; sprinkles of rain. |
| 2 | 11 47 04.4 | 58 00 | 25.3733 | 25.3808 | 644.18500 | do. do. do. |
| 3 | 12 36 05.6 | 60 00 | 20.6160 | 20.6160 | 425.01945 | do. do. do. |
| <i>Cincinnati, March 14, 1840.</i> | | | | | | |
| No. 1 | 11 ^h 33 ^m 59 ^s .2 | 48° 5' F. | 25.6600 Min. | 25.6954 Min. | 660.25358 | Beginning to rain. |
| 2 | 12 16 03.2 | 45 0 | 25.3600 | 25.4151 | 645.92731 | Raining. |
| 3 | 12 54 59.2 | 45 5 | 20.6066 | 20.6239 | 425.34525 | do. |
| <i>Louisville, Ky., March 10, 1840. Latitude 38° 03' N.; Longitude 85° 30' W.</i> | | | | | | |
| No. 1 | 11 ^h 44 ^m 02 ^s .4 | 59° 50' F. | 25.3333 Min. | 25.3363 Min. | 641.77777 | Clear; brisk N. W. wind. |
| 2 | 12 20 58.8 | 57 50 | 25.0200 | 25.0290 | 626.45084 | do. do. do. |
| 3 | 1 18 00.0 | 57 10 | 20.3333 | 20.3368 | 413.58540 | do. do. do. |
| <i>Louisville, March 11, 1840.</i> | | | | | | |
| No. 1 | 11 ^h 40 ^m 04 ^s .8 | 39° 5' F. | 25.3000 Min. | 25.3622 Min. | 643.24119 | Clear; wind moderate. |
| 2 | 12 14 04 | 41 0 | 24.9733 | 25.0421 | 627.10677 | do. do. do. |
| 3 | 12 48 03.2 | 43 | 20.3200 | 20.3400 | 413.71560 | do. do. do. |

was then deduced by Hansteen's formula. Similar experiments were made with the other needles, as below.

| | | | | | Min. Sec. |
|------------------|---------------------------|----------------------|-------------------|--|-----------|
| March 22d, 1840. | Needle No. 1. | Temperature 38°. | 500 vibrations in | | 25 34.8 |
| | | Temperature 91.4. | “ “ | | 25 44.8 |
| | | Coefficient deduced, | | | 0.000117 |
| March 25. | Mean temperature 37° 36'. | Needle No. 1. | 500 vibrations in | | 25 33.8 |
| | Mean temperature 91 5 | “ “ | “ “ | | 25 44 |
| | | Coefficient deduced, | | | 0.000122 |
| | | Mean for No. 1, | | | 0.000125 |
| March 24. | Mean temperature 50. | Needle No. 2. | 500 vibrations in | | 25 18.8 |
| | 86.12 | “ “ | “ “ | | 25 26.8 |
| | | Coefficient deduced, | | | 0.000145 |
| March 25. | Mean Temperature 48. | Needle No 3. | 500 vibrations in | | 20 34.6 |
| | 90 | “ “ | “ “ | | 20 37.6 |
| | | Coefficient deduced, | | | 0.000058 |

This last needle, the flat one with lozenge points before referred to, seems very little affected by temperature.

It may, perhaps, excite surprise, and, possibly, some doubt as to the accuracy of my observations, that the dip given above for Cincinnati is so much less than that which I obtained in November, 1837, the latter being $70^{\circ} 45'.75$, and the former $70^{\circ} 25'.5312$; and that the dip at Louisville, August 31, 1839, should have been $70^{\circ} 08'$, and on March 11, 1840, only $69^{\circ} 54'.4375$, or, as a mean, $69^{\circ} 54'.8750$. It is probable, from an experiment recorded in Major Long's expedition, making the dip at St. Louis, in 1819, $70^{\circ} 30'$, and, by my own observations, in 1839, making it, at the same place, only $69^{\circ} 31'.4$, the interval being twenty years, that the dip at this part of the earth is diminishing at the rate of about 3' per annum. This will account for part of the change observed at Cincinnati. The remainder is probably produced by the seasons, which Hansteen allows to have an influence to the extent of 15'. Allowing 7' for the annual diminution of the dip since November, 1837, we have for the change produced by the seasons,

$$\begin{aligned}
 70^{\circ} 25'.5312 + 7' + x &= 70^{\circ} 45'.75, \text{ and} \\
 x &= 70^{\circ} 45'.75 - 70^{\circ} 25'.5312 - 7' \text{ or} \\
 x &= 13' 288 = \text{the above named change.}
 \end{aligned}$$